

細胞健檢

弘晉**免費**提供**超過200種黴漿菌**檢測



你知道看不見的黴漿菌汙染
會影響實驗DATA嗎?



檢測辦法

信件主旨：InvivoGen Plasmocin™ 黴漿菌剋星 [您的學校 - 實驗室單位]



Mail 到 info@hongjing.com.tw



信件內容：
(1)我要參加細胞健檢活動
(2)實驗室聯繫電話(分機)



(1)細胞培養至少48-72小時·
細胞有8成滿以上
(2)取500 ul細胞培養液至
1.5 ml Eppendorf, 儲存於-20°C



業務專員
將在收到信件後與您聯繫
做進一步的說明

或直接撥打弘晉免付費專線：0800-008678

看不見 不代表不存在！



Mycoplasma contamination in cell cultures!

根據統計，不同地區不同實驗室約有10-85%細胞有黴漿菌污染，當細胞受到黴漿菌污染，會導致實驗結果不穩定。然而，肉眼無法判斷細胞是否遭受到黴漿菌污染且一般細胞培養用抗生素PSA，並無法防止黴漿菌污染！



弘晉提供  **InvivoGen** 全系列商品
有效 偵測 消滅 預防 黴漿菌污染



黴漿菌偵測

PlasmoTest™ - Mycoplasma Detection Kit
貨號：rep-pt1



黴漿菌消滅



Plasmocin™ treatment
貨號：ant-mpt



黴漿菌預防

Plasmocin™ prophylactic
貨號：ant-mpp

市售黴漿菌去除試劑比較

Product	Supplier	Treatment	Ease of use	Efficacy	Cytotoxicity	Resistance
BM-Cyclin	Roche	3 weeks	-	++	+	+/-
Ciprobay	Bayer	12 to 20 days	+	+	+/-	+
MRA	ICN	1 to 2 weeks	+	+	+/-	+
 Plasmocin™	InvivoGen	2 weeks	+	+++	+/-	-
 Plasmocure™	InvivoGen	2 weeks	+	++++	+	--

*2019 Sci Rep. DOI: 10.1038/s41598-018-38004-5
Metformin enhances the radiosensitizing effect of cisplatin in non-small cell lung cancer cell lines with different cisplatin sensitivities.
Riaz M.A. et al.

*2019 Sci Rep. DOI: 10.1038/s41598-018-37028-1
Neuron/Glial Antigen 2-Type VI Collagen Interactions During Murine Temporomandibular Joint Osteoarthritis.
Yotsuya M. et al.

*2018 Nat Commun. 9(1):936.
Direct conversion of injury-site myeloid cells to fibroblast-like cells of granulation tissue.
Sinha M. et al.



更多文獻請掃描



弘晉有限公司
Hong Jing Co., Ltd.

免付費專線：0800-008-678
info@hongjing.com.tw
www.hongjing.com.tw



弘晉有限公司